



Declaration of Compliance BS3882:2015

Soil source: British Sugar TOPSOIL

This declaration confirms that the topsoil represented by the attached Topsoil Analysis Report conforms to the requirements of the British Standard for Topsoil (BS3882:2015).

The sample was sampled and tested in accordance with the requirements of BS3882:2015

- Samples are taken for analysis every 8000 tonnes (5000 m3) of product
- Samples are taken from all TOPSOIL products ready for despatch
- Landscape 20 and Sports 10 are sampled after screening
- Analysis certificates are retained for a period of 5 years
- Laboratory analysis is undertaken at a **UKAS** and **MCERTS** accredited laboratory
- All laboratory methods are in accordance with BS3882:2015
- All British Sugar TOPSOIL products are produced to a **Quality Management System** approved by Lloyd's Register Quality Assurance to **ISO 9001:2008** standard

Signed

Andy Spetch

AW forth

British Sugar TOPSOIL, National TOPSOIL Manager

Sugar Way, Peterborough, PE2 9AY

Telephone 0870 2402314



Mr Andy Spetch British Sugar plc Co-Products Oundle Road Peterborough PE2 9QU

> 24th August 2021 Our Ref: TOHA/21/7117/SS Your Ref: PO 60161859

Dear Sirs

Topsoil Analysis Report: Landscape 20 Wissington

We have completed the analysis of the LANDSCAPE 20 TOPSOIL sample recently submitted, referenced *Wi-L20-Aug 21* and have pleasure reporting our findings.

The purpose of the analysis was to determine the suitability of the LANDSCAPE 20 TOPSOIL sample for general landscape purposes. In addition, this sample has been assessed to determine its compliance with the requirements of the British Standard for Topsoil (BS3882:2015 - Specification for topsoil – Table 1, Multipurpose Topsoil).

This report presents the results of analysis for the sample submitted to our office, and it should be considered 'indicative' of the topsoil source. The report and results should therefore not be used by third parties as a means of verification or validation testing or waste designation purposes, especially after the topsoil has left the British Sugar factory.

SAMPLE EXAMINATION

The sample was described as a dark greyish brown (Munsell Colour 10YR 4/2), slightly moist, friable, slightly calcareous SANDY LOAM with a weakly developed, fine to occasionally medium granular structure*. The sample was slightly stony and no unusual odours, deleterious materials, roots or rhizomes of pernicious weeds were observed.

*This appraisal of soil structure was made from examination of a disturbed sample. Structure is a key soil characteristic that may only be accurately assessed by examination in an in-situ state.

ANALYTICAL SCHEDULE

The sample was submitted to a UKAS and MCERTS accredited laboratory for a range of physical and chemical tests to confirm the composition and fertility of the soil, and the concentration of selected potential contaminants. The following parameters were determined:

- · particle size analysis;
- stone content (2-20mm, 20-50mm, >50mm);
- pH and electrical conductivity values;
- exchangeable sodium percentage;
- major plant nutrients (N, P, K, Mg);
- · organic matter content;
- C:N ratio;
- heavy metals (As, B, Ba, Be, Cd, Cr, Cu, Pb, Hg, Ni, Se, V, Zn);
- total cyanide and total (mono) phenols;
- speciated PAHs (US EPA16 suite);
- aromatic and aliphatic TPH (C5-C35 banding);
- benzene, toluene, ethylbenzene, xylene (BTEX);
- · asbestos screen.

The results are presented on the attached Certificate of Analysis and an interpretation of the results is given below. The interpretation considers the use of the LANDSCAPE 20 TOPSOIL for general landscape purposes and its compliance/non-compliance with our general landscape specification.

RESULTS OF ANALYSIS

Particle Size Analysis and Stone Content

The sample fell into the *sandy loam* texture class, which is usually considered suitable for general landscape applications provided the soil's physical condition is satisfactory.

The stone content of the sample was low and, as such, stones should not restrict the use of the soil for general landscape purposes.

pH and Electrical Conductivity Values

The sample was strongly alkaline in reaction (pH 8.3). This pH value would be considered suitable for general landscape purposes providing species with a wide pH tolerance or those known to prefer alkaline soils are selected for planting, turfing and seeding.

The electrical conductivity (salinity) value (water extract) was moderate, which indicates that soluble salts should not be present at levels that would be harmful to plants.

The electrical conductivity value by CaSO₄ extract (BS3882 requirement) fell below the maximum specified value (3300 μS/cm) given in BS3882:2015 – Table 1.

Organic Matter and Fertility Status

The sample was adequately to well supplied with organic matter and all major plant nutrients.

The C:N ratio of the sample was acceptable for general landscape purposes.

TOHA/21/7117/SS/Aug Page 2

Potential Contaminants

With reference to BS3882:2015 - Table 1: Notes 3 and 4, there is a recommendation to confirm levels of potential contaminants in relation to the topsoil's proposed end use. This includes human health, environmental protection and metals considered toxic to plants. In the absence of site-specific criteria, the concentrations that affect human health have been compared with the residential with homegrown produce land use in the Suitable For Use Levels (S4ULs) presented in the LQM/CIEH S4ULs for Human Health Risk Assessment (2015) and the DEFRA SP1010: Development of Category 4 Screening Levels for Assessment of Land Affected by Contamination – Policy Companion Document (2014). The concentration of barium has been compared with the residential land use given in the document EIC/AGS/CL:AIRE Soil Generic Assessment Criteria for Human Health Risk Assessment (2010).

Of the potential contaminants determined, none was found at levels that exceeded their guideline values.

Phytotoxic Contaminants

Of the phytotoxic (toxic to plants) contaminants determined (copper, nickel, zinc), none was found at levels that exceeded the maximum permissible levels specified in *BS3882:2015 – Table 1*.

CONCLUSION

The purpose of the analysis was to determine the suitability of the LANDSCAPE 20 TOPSOIL sample for general landscape purposes. In addition, this sample has been assessed to determine its compliance with the requirements of the British Standard for Topsoil (BS3882:2015 - Specification for topsoil – Table 1, Multipurpose Topsoil).

From the soil examination and laboratory analysis, the sample was described as a strongly alkaline, non-saline, slightly calcareous sandy loam with a weak structure and low stone content. The sample was adequately to well supplied with organic matter and all major plant nutrients. Of the potential contaminants determined, none exceeded their respective guideline values.

To conclude, based on our findings, the topsoil represented by this sample would be considered suitable for general landscape purposes (trees, shrubs and amenity grass), provided species with a wide pH tolerance or those known to prefer alkaline soils are selected and the physical condition of the soil is satisfactory.

The topsoil was also fully compliant with the requirements of the British Standard for Topsoil (BS3882:2015 – Specification for topsoil – Table 1, Multipurpose Topsoil).

RECOMMENDATIONS

Soil Handling Recommendations

It is important to maintain the physical condition of the soil and avoid structural damage during all phases of soil handling (e.g. stockpiling, respreading, cultivating, planting, seeding or turfing). As a consequence, soil handling operations should be carried out when soil is reasonably dry and non-plastic (friable) in consistency.

It is important to ensure that the soil is not unnecessarily compacted by trampling or trafficking by site machinery, and soil handling should be stopped during and after heavy rainfall and not continued until the soil is friable in consistency. If the soil is structurally damaged and compacted at any stage during the course of soiling or landscaping works, it should be cultivated appropriately to relieve the compaction and to restore the soil's structure prior to any planting, turfing or seeding.

Further details on soil handling are provided in Annex A of BS3882:2015.

TOHA/21/7117/SS/Aug Page 3

We hope this report meets with your approval and provides the necessary information. Please do not hesitate to contact the undersigned if we can be of further assistance.

Yours sincerely

Tilly Kimble-Wilde

BSc MSc

Graduate Soil Scientist

mkimble-Wilde

Rebecca Hollands BSc MSc MISoilSci Senior Soil Scientist

For & on behalf of Tim O'Hare Associates LLP

TOHA/21/7117/SS/Aug Page 4

Client:	British Sugar plc Co-Products
Client Ref:	Wissington Landscape 20
Job:	Topsoil Analysis
Date:	24/08/2021
Job Ref No:	TOHA/21/7117/SS

Client:	British Sugar plc Co-Products
Client Ref:	Wissington Landscape 20
Job:	Topsoil Analysis
Date:	24/08/2021
Job Ref No:	TOHA/21/7117/SS

Clay (<0.002mm)	%	U
Silt (0.002-0.063mm)	%	٦
Sand (0.063-2.0mm)	%	J
Texture Class (UK Classification)		J
Stones (2-20mm)	% DW	G
Stones (20-50mm)	% DW	G
Stones (>50mm)	% DW	G

pH Value (1:2.5 water extract)	units	כ
Electrical Conductivity (1:2.5 water extract)	uS/cm	כ
Electrical Conductivity (1:2 CaSO ₄ extract)	uS/cm	U
Exchangeable Sodium Percentage	%	U
Moisture Content	%	כ
Organic Matter (LOI)	%	U
Total Nitrogen (Dumas)	%	כ
C : N Ratio	ratio	כ
Extractable Phosphorus	mg/l	U
Extractable Potassium	mg/l	כ
Extractable Magnesium	mg/l	כ

Total Arsenic (As) mg/kg M Total Barium (Ba) mg/kg M Total Bervillium (Be) mg/kg M Total Evalium (Be) mg/kg M Total Cadmium (Cd) mg/kg M Total Cadmium (Cr) mg/kg M Total Camore (Cr) mg/kg M Total Copper (Cu) mg/kg M Total Copper (Cu) mg/kg M Total Lead (Pb) mg/kg M Total Michael (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Water Soluble Boron (B) mg/kg M Total (Zunide (CN) mg/kg M Total (Zunide (CN) mg/kg M Total (Innon) Phenols mg/kg M			
Total Beryllium (Be) mg/kg M Total Cadmium (Cd) mg/kg M Total Cadmium (Cd) mg/kg M Mrotal Chromium (Cr) mg/kg M Hexavalent Chromium (Cr VI) mg/kg M Total Copper (Cu) mg/kg M Total Lead (Pb) mg/kg M Total Mercury (Hg) mg/kg M Total Mickel (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Vanadium (V) mg/kg M M Total Selenium (Se) mg/kg M M Total Selanium (Be) mg/kg M M Total Cyanide (CN) mg/kg M M M M M M M M M M	Total Arsenic (As)	mg/kg	M
Total Cadmium (Cd) mg/kg M Total Chromium (Cr) mg/kg M Hexavalent Chromium (Cr VI) mg/kg M Total Copper (Cu) mg/kg M Total Lead (Pb) mg/kg M Total Mercury (Hg) mg/kg M Total Nickel (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Water Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Barium (Ba)	mg/kg	M
Total Chromium (Cr) mg/kg M Hexavalent Chromium (Cr VI) mg/kg M Total Copper (Cu) mg/kg M Total Lead (Pb) mg/kg M Total Mercury (Hg) mg/kg M Total Nickel (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Total Cyanide (CN) mg/kg M	Total Beryllium (Be)	mg/kg	М
Hexavalent Chromium (Cr VI) mg/kg M Total Copper (Cu) mg/kg M Total Lead (Pb) mg/kg M Total Mercury (Hg) mg/kg M Total Mickel (Ni) mg/kg M Total Steinum (Se) mg/kg M Total Vanadium (V) mg/kg M Total Vanadium (V) mg/kg M Total Scinc (Zn) mg/kg M Total Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Cadmium (Cd)	mg/kg	M
Total Copper (Cu) mg/kg M Total Lead (Pb) mg/kg M Total Mercury (Hg) mg/kg M Total Nickel (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Water Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Chromium (Cr)	mg/kg	M
Total Lead (Pb) mg/kg M Total Mercury (Hg) mg/kg M Total Nickel (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Total Cyanide (EN) mg/kg M Total Cyanide (CN) mg/kg M	Hexavalent Chromium (Cr VI)	mg/kg	M
Total Mercury (Hg) mg/kg M Total Nickel (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Water Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Copper (Cu)	mg/kg	M
Total Nickel (Ni) mg/kg M Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M mg/kg M Total Cyanide (CN) mg/kg M M Total Cyanide (CN) mg/kg M M Total Cyanide (CN) mg/kg M M M M M M M M M M	Total Lead (Pb)	mg/kg	M
Total Selenium (Se) mg/kg M Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Water Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Mercury (Hg)	mg/kg	M
Total Vanadium (V) mg/kg M Total Zinc (Zn) mg/kg M Water Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Nickel (Ni)	mg/kg	M
Total Zinc (Zn) mg/kg M Water Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Selenium (Se)	mg/kg	M
Water Soluble Boron (B) mg/kg M Total Cyanide (CN) mg/kg M	Total Vanadium (V)	mg/kg	M
Total Cyanide (CN) mg/kg M	Total Zinc (Zn)	mg/kg	М
	Water Soluble Boron (B)	mg/kg	M
Total (mono) Phenols mg/kg M	Total Cyanide (CN)	mg/kg	М
	Total (mono) Phenols	mg/kg	M

Naphthalene	mg/kg	М
Acenaphthylene	mg/kg	M
Acenaphthene	mg/kg	М
Fluorene	mg/kg	M
Phenanthrene	mg/kg	M
Anthracene	mg/kg	М
Fluoranthene	mg/kg	M
Pyrene	mg/kg	М
Benzo(a)anthracene	mg/kg	M
Chrysene	mg/kg	M
Benzo(b)fluoranthene	mg/kg	М
Benzo(k)fluoranthene	mg/kg	M
Benzo(a)pyrene	mg/kg	M
Indeno(1,2,3-cd)pyrene	mg/kg	M
Dibenzo(a,h)anthracene	mg/kg	M
Benzo(g,h,i)perylene	mg/kg	M
Total PAHs (sum USEPA16)	mg/kg	M

Aliphatic TPH (C5-C6)	mg/kg	M
Aliphatic TPH (C6-C8)	mg/kg	M
Aliphatic TPH (C8-C10)	mg/kg	M
Aliphatic TPH (C10-C12)	mg/kg	Μ
Aliphatic TPH (C12-C16)	mg/kg	Σ
Aliphatic TPH (C16-C21)	mg/kg	Δ
Aliphatic TPH (C21-C35)	mg/kg	M
Aliphatic TPH (C5-C35)	mg/kg	M
Aromatic TPH (C5-C7)	mg/kg	M
Aromatic TPH (C7-C8)	mg/kg	Μ
Aromatic TPH (C8-C10)	mg/kg	Μ
Aromatic TPH (C10-C12)	mg/kg	M
Aromatic TPH (C12-C16)	mg/kg	M
Aromatic TPH (C16-C21)	mg/kg	M
Aromatic TPH (C21-C35)	mg/kg	M
Aromatic TPH (C5-C35)	mg/kg	М

Aromatic TPH (C5-C35)	mg/kg	Μ
Benzene	mg/kg	M
Toluene	mg/kg	Μ
Ethylbenzene	mg/kg	M
p & m-xylene	mg/kg	M
o-xylene	mg/kg	M
Asbestos	ND/D	

p & m-xylene	mg/kg
o-xylene	mg/kg
Asbestos	ND/D

Wi-L20-Aug 21	
✓	
✓	
✓	
-	
✓	
✓	
✓	

TIM O'HARE ASSOCIATES SOIL & LANDSCAPE CONSULTANCY

8.3	✓
854	✓
2787	✓
6.9	✓
9.9	
3.7	✓
0.18	✓
12	✓
54	✓
774	✓
117	✓

7	✓
23	✓
0.3	✓
< 0.2	✓
9.8	✓
< 4.0	✓
14	✓
14	✓
< 0.3	✓
6.8	✓
< 1.0	✓
16	✓
34	✓
2.2	✓
< 1.0	✓
< 1.0	✓

< 0.05	✓
< 0.05	✓
< 0.05	✓
< 0.05	✓
< 0.05	✓
< 0.05	✓
< 0.05	✓
< 0.05	
< 0.05	1
	✓
< 0.05	Y Y Y Y
< 0.05 < 0.05	<i>Y Y Y</i>
< 0.05 < 0.05 < 0.05	<i>Y Y Y Y</i>
< 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05	\frac{}{}
< 0.05 < 0.05 < 0.05 < 0.05 < 0.05	V V V V

< 0.001	✓
< 0.001	✓
< 0.001	✓
< 1.0	✓
< 2.0	✓
< 8.0	✓
< 8.0	✓
< 10	✓
< 0.001	✓
< 0.001	✓
< 0.001	✓
< 1.0	✓
< 2.0	✓
< 10	✓
< 10	✓
< 10	✓

< 0.001	✓
< 0.001	✓
< 0.001	✓
< 0.001	✓
< 0.001	✓

Not-detected	✓

Visual Examination

The sample was described as a dark greyish brown (Munsell Colour 10YR 4/2), slightly moist, friable, slightly calcareous SANDY LOAM with a weakly developed, fine to occasionally medium granular structure The sample was slightly stony and no unusual odours, deleterious materials, roots or rhizomes of pernicious weeds were observed.

✓	Meets General Landscape Specification
Х	Fails General Landscape Specification
SL	SANDY LOAM Texture Class
M	MCERTS accredited method (& UKAS accredited method)
I	ISO 17025 accredited method
U	UKAS accredited method
G	GLP accredited method

This report presents the results of analysis for the sample submitted to our office, and it should be considered 'indicative' of the topsoil source. The report and results should therefore not be used by third parties as a means of verification or validation testing or waste designation purposes, especially after the topsoil has left the British Sugar factory.

mkimble-Wilde

Tilly Kimble-Wilde BSc MSc Graduate Soil Scientist

Results of analysis should be read in conjunction with the report they were issued with

The contents of this certificate shall not be reproduced without the express written permission of Tim O'Hare Associates LLP.